



# IgG Food Antibody Assessment (Serum)



46-50 Coombe Road  
New Malden  
Surrey KT3 4QF

63 Zillicoa Street  
Asheville, NC 28801 USA

Patient: **Order Number:**  
DOB: Completed:  
Sex: Received:  
MRN: Collected:

## IgG Food Antibody Results

| Dairy                                      | Vegetables                               | Fish/Shellfish                          | Nuts and Grains                           |
|--|--|---|---|
| Casein VL <input type="checkbox"/>         | Alfalfa 0 <input type="checkbox"/>       | Clam 0 <input type="checkbox"/>         | Almond 0 <input type="checkbox"/>         |
| Cheddar cheese VL <input type="checkbox"/> | Asparagus 0 <input type="checkbox"/>     | Cod VL <input type="checkbox"/>         | Buckwheat 0 <input type="checkbox"/>      |
| Cottage cheese VL <input type="checkbox"/> | Avocado 0 <input type="checkbox"/>       | Crab 1+ <input type="checkbox"/>        | Corn 0 <input type="checkbox"/>           |
| Cow's milk 1+ <input type="checkbox"/>     | Beets 0 <input type="checkbox"/>         | Lobster 0 <input type="checkbox"/>      | Corn gluten VL <input type="checkbox"/>   |
| Goat's milk 0 <input type="checkbox"/>     | Broccoli 0 <input type="checkbox"/>      | Oyster 0 <input type="checkbox"/>       | Gluten 1+ <input type="checkbox"/>        |
| Lactalbumin 3+ <input type="checkbox"/>    | Cabbage 0 <input type="checkbox"/>       | Red snapper VL <input type="checkbox"/> | Kidney bean 0 <input type="checkbox"/>    |
| Yogurt 1+ <input type="checkbox"/>         | Carrot 0 <input type="checkbox"/>        | Salmon 0 <input type="checkbox"/>       | Lentil 0 <input type="checkbox"/>         |
|  | Celery 0 <input type="checkbox"/>        | Sardine 0 <input type="checkbox"/>      | Lima bean 0 <input type="checkbox"/>      |
| <b>Fruits</b>                              | Cucumber 0 <input type="checkbox"/>      | Shrimp 0 <input type="checkbox"/>       | Oat 0 <input type="checkbox"/>            |
| Apple 0 <input type="checkbox"/>           | Garlic 1+ <input type="checkbox"/>       | Sole 0 <input type="checkbox"/>         | Peanut VL <input type="checkbox"/>        |
| Apricot 0 <input type="checkbox"/>         | Green Pepper 0 <input type="checkbox"/>  | Trout 0 <input type="checkbox"/>        | Pecan VL <input type="checkbox"/>         |
| Banana 0 <input type="checkbox"/>          | Lettuce 0 <input type="checkbox"/>       | Tuna VL <input type="checkbox"/>        | Pinto bean 0 <input type="checkbox"/>     |
| Blueberry 0 <input type="checkbox"/>       | Mushroom 1+ <input type="checkbox"/>     |   | Rice 0 <input type="checkbox"/>           |
| Cranberry 0 <input type="checkbox"/>       | Olive 0 <input type="checkbox"/>         | <b>Poultry/Meats</b>                    | Rye 0 <input type="checkbox"/>            |
| Grape 0 <input type="checkbox"/>           | Onion 0 <input type="checkbox"/>         | Beef 0 <input type="checkbox"/>         | Sesame VL <input type="checkbox"/>        |
| Grapefruit 0 <input type="checkbox"/>      | Pea 0 <input type="checkbox"/>           | Chicken 0 <input type="checkbox"/>      | Soy 0 <input type="checkbox"/>            |
| Lemon 0 <input type="checkbox"/>           | Potato, sweet 0 <input type="checkbox"/> | Egg white VL <input type="checkbox"/>   | Sunflower seed 0 <input type="checkbox"/> |
| Orange 0 <input type="checkbox"/>          | Potato, white 0 <input type="checkbox"/> | Egg yolk VL <input type="checkbox"/>    | Walnut VL <input type="checkbox"/>        |
| Papaya 0 <input type="checkbox"/>          | Spinach 0 <input type="checkbox"/>       | Lamb 0 <input type="checkbox"/>         | Wheat 1+ <input type="checkbox"/>         |
| Peach 0 <input type="checkbox"/>           | String bean 0 <input type="checkbox"/>   | Pork 0 <input type="checkbox"/>         |   |
| Pear 0 <input type="checkbox"/>            | Tomato 0 <input type="checkbox"/>        | Turkey 0 <input type="checkbox"/>       | <b>Miscellaneous</b>                      |
| Pineapple 0 <input type="checkbox"/>       | Zucchini 0 <input type="checkbox"/>      |   | Yeast 0 <input type="checkbox"/>          |
| Plum 0 <input type="checkbox"/>            |  |   | Cane sugar 0 <input type="checkbox"/>     |
| Raspberry 0 <input type="checkbox"/>       |  |   | Chocolate 0 <input type="checkbox"/>      |
| Strawberry 0 <input type="checkbox"/>      |  |   | Coffee 0 <input type="checkbox"/>         |
| <b>Total IgE</b>                           |  |   |   |
| Inside                                     |  | Outside                                 | Reference Range                           |
| Total IgE ♦ <input type="checkbox"/>       |  | 179.0                                   | <=87.0 IU/mL                              |

0  None Detected    VL  Very Low    1+  Low    2+  Moderate    3+  High

- The performance characteristics of all assays have been verified by Genova Diagnostics, Inc. Unless otherwise noted with ♦, the assay has not been cleared by the U.S. Food and Drug Administration.
- Total IgE level may have clinical significance regardless of specific antibody levels.
- Increasing levels of antibody detected suggest an increasing probability of clinical reactivity to specific foods.
- The Elimination Diet commentary is specific to IgG results only. Allergens inducing an IgE response should be completely avoided.

### Laboratory Comments

# Summary of IgG Test Results

## Reactive / Non-Reactive Foods

### 3+ High

Lactalbumin

### 1+ Low

Cow's milk  
MushroomCrab  
WheatGarlic  
Yogurt

Gluten

### VL Very Low

Casein  
Cottage cheese  
Pecan  
WalnutCheddar cheese  
Egg white  
Red SnapperCod  
Egg yolk  
SesameCorn gluten  
Peanut  
Tuna

### 0 None Detected

Alfalfa  
Asparagus  
Beets  
Cabbage  
Chicken  
Corn  
Grape  
Lamb  
Lima bean  
Onion  
Pea  
Pinto bean  
Potato, white  
Salmon  
Soy  
Sunflower seed  
YeastAlmond  
Avocado  
Blueberry  
Cane sugar  
Chocolate  
Cranberry  
Grapefruit  
Lemon  
Lobster  
Orange  
Peach  
Plum  
Raspberry  
Sardine  
Spinach  
Tomato  
ZucchiniApple  
Banana  
Broccoli  
Carrot  
Clam  
Cucumber  
Green pepper  
Lentil  
Oat  
Oyster  
Pear  
Pork  
Rice  
Shrimp  
Strawberry  
TroutApricot  
Beef  
Buckwheat  
Celery  
Coffee  
Goat's milk  
Kidney bean  
Lettuce  
Olive  
Papaya  
Pineapple  
Potato, sweet  
Rye  
Sole  
String bean  
Turkey

## Commentary

### Overview

**Immunoglobulin G (IgG) antibodies** that elicit an immune response to food are in a class distinct from Immunoglobulin E (IgE) food allergy reactions. IgG-mediated food responses are described as delayed hypersensitivity reactions and have been associated in the peer-reviewed literature with an array of common clinical conditions including migraine, obesity, asthma, autoimmune diseases, and irritable bowel syndrome.

#### **IgG Testing: Factors to Consider**

IgG testing can be very useful in screening foods that a person is eating on a regular basis and which may be causing adverse reactions. However, it is possible to have adverse reactions to foods with low or non-detected levels of IgG. Because the IgG profile measures exposure of the immune system to food antigens, performing this test on a patient who is not consuming a particular food or who is taking a drug with known ability to suppress immune function (i.e. steroids) may result in the test not showing a positive reaction, potentially leading to a false negative result for the particular food. Be advised that if the patient is already on an elimination diet due to known food reactions, a negative result on an IgG food antibody profile does not necessarily mean that they can freely eat the food without experiencing symptoms.

#### **IgG Results Interpretation**

The amount of IgG antibodies is measured using a semi-quantitative ELISA assay procedure. The relative degrees of IgG present for each food are reported using a semi-quantitative level; None Detected (0), VL (very low), Low (1+), Moderate (2+) or High (3+). The degree of reactivity may not correlate with the severity of patient's response, therefore clinical correlation is advised as it can help direct treatment.

#### **Clinical Management of Reactive IgG Foods: Elimination Diet**

The purpose of an elimination diet is to pinpoint symptom-triggering foods that may be the root cause of and/or perpetuating chronic health issues. This diet is specific to food sensitivities that elicit an Immunoglobulin G (IgG) response and not those defined as classic (IgE-mediated) food allergy reactions. An elimination diet is a strategic process which depends on the oversight of the healthcare provider to ensure that a patient's nutritional requirements - macronutrient, micronutrient, and caloric needs - are adequate.

### Four-Phases of an Elimination Diet



#### **PHASE 1 – PREPARATION**

A patient's clinical presentation and the IgG Food Antibody Assessment results typically determine which food(s) to temporarily remove from the diet. The average time frame for an elimination diet is 1 to 3 months. It is optimal to work with the patient to determine a start and end date for the elimination diet. Patient guidance around preparation ahead of the start date is important to ensure success. These include: (1) encouraging the patient to remove offending foods from the home and adjust grocery shopping accordingly; (2) providing the patient with resources that advance meal preparation, such as recipe books or reputable websites. Directing the patient to record foods consumed, date of consumption/elimination, and any notable changes in symptoms in a food journal can help track the progress of the diet.

## Commentary



### PHASE 2 – ELIMINATION

It is important to ensure the patient avoids those foods which resulted in a demonstrable reaction, either in whole food forms or as ingredients in other prepared foods to gain the greatest benefit. For patients unable to eliminate all reactive foods from their diet, focusing on the foods that elicited a stronger reaction (i.e.: 2+ and 3+) may be considered for an elimination diet. Practitioners may also encourage elimination of a complete food group when the patient shows reactivity to all foods tested within that group.



### PHASE 3 – REINTRODUCTION

The reintroduction of eliminated foods is done one food at a time while monitoring for any adverse reaction. The patient should consume the test food several times throughout the day for several days. If symptoms occur with reintroduction, the patient should be instructed to remove that food once again and to evaluate whether the symptoms diminish over the next few days following elimination. Signs which may indicate an IgG food reaction include the following: headache, itching, bloating, fatigue, diarrhea or constipation, and indigestion. If the food does not cause symptoms during the reintroduction phase, it can be added back into the diet. The patient should continue this process with each food eliminated.

**CAUTION:** *All patients warrant counseling related to signs and management of immediate hypersensitivity reactions prior to food reintroduction following an elimination diet. If reintroduction of a food causes an immediate allergic reaction (i.e. swelling of face, mouth, tongue, etc.; wheezing, rash/hives, or other allergic symptoms), it is imperative that the patient be treated as soon as possible. Following resolution of the immediate hypersensitivity reaction, the patient should be instructed to completely avoid consumption of that food.*



### PHASE 4 – LONG TERM MANAGEMENT

An elimination diet based on food sensitivity testing is part of a comprehensive approach to overall gastrointestinal health. Based on the test results and the complete clinical presentation of the patient, a long-term plan is usually developed utilizing the results of the reintroduction phase. Clinicians may also consider assessing and treating intestinal permeability, as gut barrier integrity is important for proper immune responses to foods. Nutrients that have been found to support intestinal barrier and decrease potential inflammation are glutamine, vitamin A, vitamin D, essential fatty acids (Omega-3), probiotics, and butyrate. Botanicals that can also be considered to assist with intestinal health are slippery elm, deglycyrrhizinated licorice (DGL), Aloe vera extract, and marshmallow root.



# IgE Food Antibody Assessment



46-50 Coombe Road  
New Malden  
Surrey KT3 4QF

63 Zillicoa Street  
Asheville, NC 28801 USA

Patient:  
DOB:  
Sex: F  
MRN:

Order Number:  
Completed:  
Received:  
Collected:

## IgE Food Antibody Results

|               | RESULT<br>kU/L | CLASS | INDICATOR                |                | RESULT<br>kU/L | CLASS | INDICATOR                |
|---------------|----------------|-------|--------------------------|----------------|----------------|-------|--------------------------|
| <b>Grains</b> |                |       |                          | <b>Nuts</b>    |                |       |                          |
| Buckwheat     | <0.24          | 0/1   | <input type="checkbox"/> | Almond         | 0.43           | II    | <input type="checkbox"/> |
| Corn          | <0.24          | 0/1   | <input type="checkbox"/> | Brazil nut     | <0.24          | 0/1   | <input type="checkbox"/> |
| Oat           | <0.24          | 0/1   | <input type="checkbox"/> | Coconut        | <0.24          | 0/1   | <input type="checkbox"/> |
| Rice          | <0.24          | 0/1   | <input type="checkbox"/> | Hazelnut       | 0.97           | II    | <input type="checkbox"/> |
| Sesame        | <0.24          | 0/1   | <input type="checkbox"/> | Peanut         | <0.24          | 0/1   | <input type="checkbox"/> |
| Soybean       | <0.24          | 0/1   | <input type="checkbox"/> | <b>Seafood</b> |                |       |                          |
| Wheat         | <0.24          | 0/1   | <input type="checkbox"/> | Blue mussel    | <0.24          | 0/1   | <input type="checkbox"/> |
| <b>Dairy</b>  |                |       |                          | Codfish        | <0.24          | 0/1   | <input type="checkbox"/> |
| Egg white     | <0.24          | 0/1   | <input type="checkbox"/> | Salmon         | <0.24          | 0/1   | <input type="checkbox"/> |
| Cow's milk    | <0.24          | 0/1   | <input type="checkbox"/> | Shrimp         | <0.24          | 0/1   | <input type="checkbox"/> |
|               |                |       |                          | Tuna           | <0.24          | 0/1   | <input type="checkbox"/> |

## Total IgE

|           |                               |                                    |                 |
|-----------|-------------------------------|------------------------------------|-----------------|
|           | Inside                        | Outside                            | Reference Range |
| Total IgE | <input type="text" value=""/> | <input type="text" value="179.0"/> | <=87.0 IU/mL    |

- IgE levels must be used in conjunction with the clinical picture and are not intended to be independently diagnostic.
- The performance characteristics of all assays have been verified by Genova Diagnostics, Inc. All assays are cleared by the U.S. Food and Drug Administration.
- Total IgE level may have clinical significance regardless of specific antibody levels.
- Increasing levels of antibody detected suggest an increasing clinical reactivity to specific foods.

## Key

| Class | kU/L        | Levels of Specific IgE    | Indicator                |
|-------|-------------|---------------------------|--------------------------|
| 0/1   | <=0.24      | Undetectable or Equivocal | <input type="checkbox"/> |
| I     | 0.25 - 0.39 | Low                       | <input type="checkbox"/> |
| II    | 0.4 - 1.29  | Moderate                  | <input type="checkbox"/> |
| III   | 1.3 - 3.89  | High                      | <input type="checkbox"/> |
| IV    | 3.9 - 14.99 | Very High                 | <input type="checkbox"/> |
| V     | 15 - 24.99  | Very High                 | <input type="checkbox"/> |
| VI    | >=25        | Very High                 | <input type="checkbox"/> |

## Laboratory Comments